



E693
JACC April 5, 2011
Volume 57, Issue 14



IMAGING AND DIAGNOSTIC TESTING

PREDICTION OF CARDIAC RESYNCHRONIZATION THERAPY RESPONSE - VALUE OF CARDIAC 123-IODINE META-IODOBENZYLGUANIDINE IMAGING

ACC Poster Contributions

Ernest N. Morial Convention Center, Hall F

Sunday, April 03, 2011, 10:00 a.m.-11:15 a.m.

Session Title: Radionuclide Imaging: New Applications

Abstract Category: 39. Nuclear Cardiology/PET

Session-Poster Board Number: 1027-184

Authors: *See Hooi Ewe, Mark J. Boogers, Petra Dibbets-Schneider, Rutger J. van Bommel, Matteo Bertini, Victoria Delgado, Arnold C.T. Ng, Dominique Auger, Tomasz G. Witkowski, Imad Al Younis, Bernies van der Hiel, Martin J. Schalij, Ernst E. van der Wall, Jeroen J. Bax, Leiden University Medical Center, Leiden, The Netherlands*

Background: It is unclear whether sympathetic abnormality, as assessed with cardiac 123-iodine meta-iodobenzylguanidine (123I-mIBG) imaging, can influence left ventricular (LV) reverse remodeling post cardiac resynchronization therapy (CRT). This study aimed to evaluate the usefulness of cardiac 123I-mIBG parameters in predicting CRT response.

Methods: A total of 101 heart failure patients referred for CRT were included. All patients underwent cardiac 123I-mIBG imaging prior to CRT. Heart-to-mediastinum ratio (HMR) and myocardial washout rate were obtained from planar imaging while SPECT imaging was performed to obtain summed 123I-mIBG SPECT defect score. Baseline echocardiography was performed including the assessment of LV dyssynchrony. At 6 months, echocardiography was repeated and CRT response was defined as a reduction of >15% in LV end-systolic volume.

Results: Fifty-six patients showed CRT response at 6 months. Responders were less likely to have ischemic etiology (61% vs. 87%, $p=0.006$) but more extensive LV dyssynchrony (74 ± 43 ms vs. 54 ± 41 ms, $p=0.03$) at baseline. On 123I-mIBG imaging, CRT responders had higher early (1.59 ± 0.21 vs. 1.48 ± 0.18 , $p=0.009$) and late HMR (1.48 ± 0.20 vs. 1.33 ± 0.16 , $p<0.001$), lower washout rate ($6.6\pm 5.5\%$ vs. $10.0\pm 6.6\%$, $p=0.008$) and lower early summed SPECT defect score (20.4 ± 8.9 vs. 25.6 ± 9.9 , $p=0.01$). On multiple logistic regression analysis, every 0.1 unit increase in late HMR was independently associated with response to CRT (OR=1.49, $p=0.013$). Importantly, the combination of late HMR and LV dyssynchrony resulted in a significantly improved prediction of CRT response over LV dyssynchrony alone, as the c-statistic increased from 0.63 to 0.75 ($p=0.044$).

Conclusions: Baseline late HMR on 123I-mIBG planar imaging has an incremental value, in addition to baseline LV dyssynchrony, for prediction of response to CRT in patients with advanced heart failure.